Market Assets

- Underlyings: stocks, bonds, foreign exchange
- \cdot Derivatives
 - forward and futures contracts
 - vanilla options (puts and calls)
 - exotic options

Stocks

Share of ownership of tangible company ("equity") Value depends on future earnings prospects at least market's opinion of what other people's opinion will be ("irrational exuberance" ...)

Efficient market hypothesis:

Current price encodes all information available anywhere. (If everyone thinks it will go up, it will already be up.) Price changes depend on arrival of unanticipated information

 \rightarrow suggests random walk models

We can anticipate *magnitude* of future movements, not *sign*. Past history is good guide to *statistical properties* of future movements (unless option trading modifies properties).

Bonds

Committment to pay specific amount of cash

at specific future time *T*

Issued by corporations (default risk) and government (no risk) Present value depends on interest rate r between now and T. r constant and known: present value $B_0 = e^{-rT}B_T$

E.g. overnight rate between large corporations ("repo rate") (Short-term option pricing not too sensitive to r.)

Interesting issues:

- Dynamics of interest rates for different times (*yield curve*).
 Interest rate fluctutates "randomly", like equities but many more "degrees of freedom". Mathematically challenging.
- *Credit spreads:* additional interest demanded for borrower risk of default. Big events, small probability.

Forward and futures contracts Agreement to buy asset for specific price at specific time. No choice for either party.


- strike (delivery) price : K
 - expiration date: *T*
 - value of contract: *F*
- market price of underlying asset: *S*

At t = T, price of contract in terms of price of underlying:

$$F_T = S_T - K$$

Often just exchange this amount of cash. Futures: standardized and traded on exchange

Payoff function of futures contract:



Determine present value F_0 by no-arbitrage reasoning: You promise to deliver one ounce gold time T in future for K. You have risk that gold might go up between now and T. Solution: go buy gold now, invest difference at risk-free rate.

Purchase one ounce gold now: cost S_0

Borrow Ke^{-rT}

Net cash from pocket: $S_0 - Ke^{-rT}$

At expiry:

Customer gives you delivery price *K* You pay off the loan (loan amount has grown to *K*) You give him the gold You are clear The *only* proper price to charge for this contract is the cost of setting up the hedging portfolio:

$$F_0 = S_0 - K e^{-rT}$$


- If contract is being bought for more than this, then sell a contract, buy gold and borrow cash as above, and pocket a riskless profit.
- If contract is being sold for less than this, then buy a contract, sell short one ounce of gold, and loan out the cash difference. At time *T* you have a riskless profit.

Delivery price $K = e^{rT}S_0$ often set so $F_0 = 0$ Futures price is proxy for current price (often more liquid) Currency futures: include foreign interest Options Buyer has the *right* but not the *obligation* to execute the deal

> Call option: right to purchase asset for price *K* Put option: right to sell asset for price *K*

Buy or sell to whoever holds the other side. Value of option at expiry in terms of underlying

> Call: $C_T = \max\{S_T - K, 0\}$ Put: $P_T = \max\{K - S_T, 0\}.$



Long side: never negative because no obligation corners are convex because choose better value

Long call, short put: *bullish* exposure (make money if price ↗) Long put, short call: *bearish* exposure (make money if price ↘)

Speculate: Use options to enhance risk and reward (?) *Hedge:* Use options to decrease exposure to preexisting risk

Evaluate option value using no-arbitrage pricing Harder than forwards because payoff is nonlinear Major distinction: *European:* Can exercise *only* at expiry time Analytic expression for solution *American:* Can exercise anytime up to expiry (most are this). Optimal control/decision problem Gives PDE with free boundary (hard)


Various types of exotic options: *Spreads, straddles, etc:* superpositions of vanilla any payoff function V(S, T) *Bermudan:* Exercise at finitely many specified times *Asian:* Payoff based on time average of asset value *Lookback:* Payoff based on *max* or *min* of value *Barrier:* Become worthless/valuable if price crosses a level Put/call parity: Consider European call option European put option futures contract all on same asset with same strike price KExamine payoff: at t = T,

$$C - P = F.$$

Therefore also true at earlier times:

$$C = P + S - Ke^{-rT}$$

Can get either call or put value from the other Useful to decompose more complicated instruments



Continuum limit

Above procedure valid for any *N*, any set of node prices Real world has (more or less) continuous time continuous price evolution (actually, "tick" size)

Take $N \rightarrow \infty$ for continuous time sampling Take node spacing $\rightarrow 0$ for continuous price sampling

If model is reasonable, solution should converge to something independent of details of tree


Discrete formula $V_0 = e^{-rT}(qV_2 + (1 - q)V_2)$ will converge to *Black-Scholes partial differential equation* for V(S, t)Solve PDE with known values V(S, T), extract $V(S_0, 0)$ Need to specify tree more systematically Choose tree to have constant up/down *ratios* at each node Children of $S_{i,j}$ are $S_{i+1,j}$ and $S_{i+1,j+1}$

$$S_{i+1,j+1} = u S_{i,j} \qquad S_{i+1,j} = d S_{i,j}$$

u, d uniform across tree, but depend on N

Proportional increments rather than fixed arithmetic jumps:

- maybe more reasonable:
 size of motion should scale with value of price
- mathematical problems if price becomes negative (might not happen if *T* is small)



Time levels equally spaced: k = T/NTwo degrees of freedom:

$$u = \exp\left[sk + \frac{1}{2}h\right]$$
 $d = \exp\left[sk - \frac{1}{2}h\right]$.

"spread" h:
$$\frac{u-d}{d} = e^h - 1 \approx h$$

"drift" s: $\sqrt{ud} = e^{sk}$

Want to take $h \rightarrow 0$ and *s* constant as $k \rightarrow 0$.

- Value of *s* will not matter (compensated by *q*)
- Relationship of h to k will be critical

Option value depends on some properties of tree but not others

Pricing on tree: same strategy as before

risk-neutral probabilities

$$q = \frac{e^{rk} - d}{u - d} = \frac{e^{(r-s)k} - e^{-h/2}}{e^{h/2} - e^{-h/2}}$$

inequality constraint

$$-\frac{1}{2}h < (r-s)k < \frac{1}{2}h.$$

iteration formula for option value

$$V_{i,j} = e^{-rk} \left(q V_{i+1,j+1} + (1-q) V_{i+1,j} \right)$$

Values V(S, T) given at t = T: starting values

$$V_{N,j} = V(S_{N,j}, N)$$

To take limit, need to specify relationship h to kTake

$$k = \lambda h^2$$
, λ constant as $h, k \to 0$

or

$$h = \sqrt{k/\lambda}$$

How do we know that $k \sim h^2$ is right scaling? To confirm, check final result in limit $\lambda \to 0, \infty$.

(Note: inequality constraint always satisfied when *h* small.)

Aside: convergence of finite-difference method to PDE in simple problem

Consider staggered grid of points

$$t_i = i k, \quad i = 0, 1, \dots$$

 $x_{i,j} = \left(\frac{1}{2}i + j\right) h, \quad -\infty < j < \infty$

Point (i, j) has "ancestors" (i - 1, j), (i - 1, j + 1)Define $u_{i,j}$ by

1.
$$u_{0,j} = f(jh)$$
, smooth function $f(x)$
2. $u_{i,j} = \frac{1}{2} (u_{i-1,j} + u_{i-1,j+1})$

Theorem: As $h, k \to 0$ with $k = \lambda h^2$, then $u_{i,j} \to u(x_{i,j}, t_i)$, where u(x, t) is solution of initial-value PDE

$$u(x,0) = f(x)$$
$$\frac{\partial u}{\partial t} = \frac{1}{4\lambda} \frac{\partial^2 u}{\partial x^2}$$

Proof: Based on *Lax equivalence theorem* for finite-difference schemes

Consistency + Stability \Rightarrow Convergence

Stability: obvious since $u_{i,\cdot}$ is smoothing of $u_{i-1,\cdot}$

Consistency: replace $u_{i,j} = u(x_{i,j}, t_i)$ exactly, with $u_t = \frac{1}{4\lambda}u_{xx}$ see by how much difference formula fails to be satisfied:

Make local expansions about point $(i - 1, j + \frac{1}{2})$.

$$u_{i,j} \sim u + ku_t + \cdots$$

$$u_{i-1,j} \sim u - \frac{h}{2}u_x + \frac{1}{2}\left(\frac{h}{2}\right)^2 u_{xx} + \cdots$$

$$u_{i-1,j+1} \sim u + \frac{h}{2}u_x + \frac{1}{2}\left(\frac{h}{2}\right)^2 u_{xx} + \cdots$$

$$u_{i,j} - \frac{1}{2} \left(u_{i-1,j} + u_{i-1,j+1} \right) \sim k u_t - \left(\frac{h}{2} \right)^2 u_{xx} + \cdots$$
$$\sim k \left(u_t - \frac{1}{4\lambda} u_{xx} \right) + \cdots$$

So consistency requires

$$u_t = \frac{1}{4\lambda} u_{XX}$$

QED

Not to hard to make this fully rigorous

end of Aside

Option-pricing problem:

Assume smooth solution, make local asymptotic expansions about suitable intermediate point

$$V_{i,j} = V - kv_t + \frac{1}{2}k^2V_{tt} + \cdots$$
$$V_{i+1,j} = V - h_-V_S + \frac{1}{2}h_-^2V_{SS} + \cdots$$
$$V_{i+1,j+1} = V + h_+V_S + \frac{1}{2}h_+^2V_{SS} + \cdots$$

Compute consistency just as for simple model problem

The Black-Scholes equation for derivative pricing:

$$V_t + \frac{1}{2}\sigma^2 S^2 V_{SS} + r S V_S - r V = 0.$$

$$\frac{1}{2}\sigma^2 = \frac{1}{8\lambda}$$

Satisfied by value of *every* derivative security on *S*. (As long as no early exercise or other complication)

- Backwards parabolic: *terminal* data at t = T
- Linear: asset values add

 σ^2 appears as a diffusion coefficient related to how much stock price "jumps around"

s has vanished! σ tells us how to construct grid: $h = 2\sigma\sqrt{k}$

How do we know σ ?

How do we know other grids might not give different PDE? (stochastic calculus helps here)

Solution of Black-Scholes equation

new independent variables

$$x = \log \frac{S}{S_{\text{ref}}}, \quad \tau = \sigma^2 (T-t)$$

au is time remaining to expiration

New dependent function

$$V(S,t) = V_{\text{ref}} u \left(\log \frac{S}{S_{\text{ref}}}, \sigma^2(T-t) \right)$$

Get

$$u_{\tau} = \frac{1}{2}u_{xx} + \beta u_x - \gamma u,$$

with constant coefficients

$$\beta = \frac{r - \frac{1}{2}\sigma^2}{\sigma^2}, \qquad \gamma = \frac{r}{\sigma^2}.$$

Easy to solve initial-value problem using Green's functions Black-Scholes formula for European call with strike *K*:


$$C(S,t) = SN\left(\frac{\log(S/K) + \left(r + \frac{1}{2}\sigma^2\right)(T-t)}{\sigma\sqrt{T-t}}\right)$$
$$- Ke^{-r(T-t)}N\left(\frac{\log(S/K) + \left(r - \frac{1}{2}\sigma^2\right)(T-t)}{\sigma\sqrt{T-t}}\right)$$

Get European put by put-call parity

Compare with market-quoted option values Impossible to find single volatility value that works for all strikes (even at just one maturity date)

Historical volatility: The volatility estimated from statistical analysis of past price motions.

Implied volatility: The volatility you have to use in the Black-Scholes pricing formula to get the market price.



Dynamic Hedging:

$$\Delta = \frac{V_2 - V_1}{S_2 - S_1} \rightarrow \frac{\partial V}{\partial S}$$

How much you have to hold at each instant to maintain hedge Jumps around as much as stock price

Vanilla puts and calls as $t \nearrow T$:

- $\Delta \rightarrow \pm 1$ if option is expiring in the money
- $\Delta \rightarrow 0$ if expiring out of the money

You end with exactly what you need

American option At each moment, a *decision problem:* Should I exercise, or should I hold??

Option value always \geq value of exercise (e.g. $V(S,t) \geq \Lambda(S,t) = \max\{S - K, 0\}$ for American call)

Tree modification for early exercise:

Compare computed value at each node against exercise value Use the larger of the two.

This affects values at all nodes *earlier* in time.

Black-Scholes equation with early exercise:

$$-V_t \ge \frac{1}{2}\sigma^2 S^2 V_{SS} + \gamma S V_S - \gamma V$$
 and $V(S,t) \ge \Lambda(S,t)$

And at least one of them is an equality

A variational inequality, or obstacle problem

Free boundary problem:

critical stock price $S_*(t)$ where you should exercise American put: exercise for $S < S_*$ American call: exercise for $S > S_*$, if stock pays dividends.

Physical analogs: oxygen consumption problem, Stefan problem

Stochastic models Binomial tree ⇒ random walk Assign probabilities to up and down moves These probabilities disappear from final solution

Stock motion has probability distribution Changes in $\log S(t)$ are serially independent

$$\log \frac{S(t)}{S(0)}$$
 is Gaussian

mean = μt variance = $\sigma^2 t$

Model from Bachelier (1900) (for arithmetic motion) First mathematics of Brownian motion (before Einstein) Stochastic differential equation

 $dS = \mu S dt + \sigma S dX,$

dX is increment of standard Brownian

Theory is elegant, obscures messy reality