Lecture 2: Fitting the Volatility Skew

Jim Gatheral, Merrill Lynch*

Case Studies in Financial Modelling Course Notes, Courant Institute of Mathematical Sciences, Fall Term, 2001

^{*}I am grateful to Peter Friz for carefully reading these notes, providing corrections and suggesting useful improvements.

5 Getting Implied Volatility from Local Volatilities

5.1 Motivation

For a model to be useful in practice, it needs to return (at least approximately) the current market prices of European options. That implies that we need to fit the parameters of our model (whether stochastic or local volatility model) to market implied volatilities. A number of ways have been suggested to do this for local volatility models. For example, we could work with the European option prices directly in a trinomial tree framework as in Derman, Kani, and Chriss (1996) or we could maximize relative entropy (of missing information) as in Avellaneda, Friedman, Holmes, and Samperi (1997). These methods are non-parametric (assuming actual option prices are used, not interpolated or extrapolated values). On the other hand, we could parameterize the risk-neutral distributions as in Rubinstein (1998) or parameterize the implied volatility surface directly as in Shimko (1993). For a recent review of the literature, see Jackwerth (2000).

It might be surprising at first to learn that getting local volatilities from the implied volatility surface is very difficult in practice given that we have a reasonably straightforward formula for doing that. The problem is that we don't have a complete implied volatility surface, we only have a few bids and offers per expiration. To apply a parametric method, we need to interpolate and extrapolate the known implied volatilities. It is very difficult to do this without introducing arbitrage. The arbitrages to avoid are roughly speaking, negative vertical spreads, negative butterflies and negative calendar spreads (where the latter are carefully defined). Even non-parametric methods fail because of noise in the prices and the bid/offer spread.

In what follows, we will concentrate on the implied volatility structure of stochastic volatility models so we won't have to worry about the possibility of arbitrage which is excluded from the outset.

First, we derive an expression for implied volatility in terms of local volatilities. In principle, this should allow us to investigate the shape of the implied volatility surface for any local volatility or stochastic volatility model because we know from Section 2.5 how to express local variance as an expectation of instantaneous variance in a stochastic volatility model.

5.2 A Formula to get Implied Volatility from Local Volatility

In Section 2.3, we saw how to get local volatilities from implied volatilities. We could try to invert the complicated-looking equation for local volatility in terms of implied given in that section. However, it is no surprise that this approach doesn't yield any easy results (at least not to me).

Instead, by extending the work of Blacher (1998), we derive a general path-integral representation of Black-Scholes implied variance. We start by assuming that the stock price S_t satisfies the (local volatility) SDE

$$\frac{dS_t}{S_t} = \mu_t dt + \sigma_{t,S_t} dZ_t$$

and that the market prices contingent claims accordingly so that, in particular, the value V of a contingent claims must satisfy a generalization of the Black-Scholes equation:

$$\frac{1}{2}\sigma_{t,S_{t}}^{2}S_{t}^{2}\frac{\partial^{2}V}{\partial S_{t}^{2}} + \mu S_{t}\frac{\partial V}{\partial S_{t}} = -\frac{\partial V}{\partial t}$$

Path-by-path, for any suitably smooth function $f(S_t, t)$ of the random stock price S_t , the difference between the initial value and the final value of the function $f(S_t, t)$ is obtained by anti-differentiation. Then, applying Itô's Lemma, we get

$$f(S_T, T) - f(S_0, 0) = \int_0^T df$$

=
$$\int_0^T \left\{ \frac{\partial f}{\partial S_t} dS_t + \frac{\partial f}{\partial t} dt + \frac{\sigma_{S_t, t}^2 S_t^2}{2} \frac{\partial^2 f}{\partial S_t^2} dt \right\} (23)$$

In particular, the Black-Scholes (BS) formula $C_{BS}(S_t, t, T)$ for a call option expiring at time T with some arbitrary time-dependent volatility parameter is a smooth function of the stock price and must satisfy equation (23). Recall the form of the Black-Scholes formula

$$C_{BS}(S_t, K, t, T) = F_{t,T} N(d_1) - K N(d_2)$$

with

$$d_1 = \frac{\ln(F_{t,T}/K)}{\sigma_{BS}\sqrt{T}} + \frac{\sigma_{BS}\sqrt{T}}{2}; \ d_2 = d_1 - \sigma_{BS}\sqrt{T}$$

where the time -T forward price at time t is denoted by $F_{t,T}$, the strike price of the option by K and $\sigma_{BS} = \sigma_{BS}(K, t, T)$ is the Black-Scholes implied volatility which is of course a function of calendar time t, strike K and T.

 $C_{BS}(S_t, K, t, T)$ must satisfy the Black-Scholes equation (assuming zero interest rates and dividends):

$$\frac{\partial C_{BS}}{\partial t} = -\frac{1}{2} v_{K,T}(t) S_t^2 \frac{\partial^2 C_{BS}}{\partial S_t^2}$$
(24)

where the forward Black-Scholes variance $v_{K,T}(s)$ is given by

$$v_{K,T}(s) = \frac{\partial}{\partial s} \left\{ \sigma_{BS}(K, t, s)^2(s - t) \right\}$$

Under the usual assumptions, the non-discounted value $C(S_0, K, 0, T)$ of a call option is given by the expectation of the final payoff under the risk-neutral measure. Then, applying (23), we obtain:

$$C(S_{0}, K, 0, T) = \mathbf{E} \left[(S_{T} - K)^{+} | S_{0} \right]$$

$$= \mathbf{E} \left[C_{BS} \left(S_{T}, K, T, T \right) | S_{0} \right]$$

$$= C_{BS} \left(S_{0}, K, 0, T \right) + \mathbf{E} \left[\int_{0}^{T} \left\{ \frac{\partial C_{BS}}{\partial S_{t}} dS_{t} + \frac{\partial C_{BS}}{\partial t} dt + \frac{1}{2} \sigma_{S_{t}, t}^{2} S_{t}^{2} \frac{\partial^{2} C_{BS}}{\partial S_{t}^{2}} dt \right\} \middle| S_{0} \right]$$

(25)

Finally, we use the BS equation to substitute for the time derivative $\frac{\partial C_{BS}}{\partial t}$ and obtain:

$$C(S_{0}, K, 0, T) = C_{BS}(S_{0}, K, 0, T) + \mathbf{E} \left[\int_{0}^{T} \left\{ \frac{\partial C_{BS}}{\partial S_{t}} dS_{t} + \frac{1}{2} \left\{ \sigma_{S_{t},t}^{2} - v_{K,T}(t) \right\} S_{t}^{2} \frac{\partial^{2} C_{BS}}{\partial S_{t}^{2}} dt \right\} \middle| S_{0} \right]$$

$$= C_{BS}(S_{0}, K, 0, T) + \mathbf{E} \left[\int_{0}^{T} \frac{1}{2} \left\{ \sigma_{S_{t},t}^{2} - v_{K,T}(t) \right\} S_{t}^{2} \frac{\partial^{2} C_{BS}}{\partial S_{t}^{2}} dt \middle| S_{0} \right]$$
(26)

where the second equality uses the fact that S_t is a martingale.

By definition of implied volatility, $C_{BS}(S_0, K, 0, T) = C(S_0, K, 0, T)$ when $v_{K,T}$ is the Black-Scholes implied forward variance (*i.e.* the Black-Scholes formula must give the market price of the option). Then the second term in equation (26) must vanish. A sufficient condition for this is to have

$$v_{K,T}(t) = \frac{\mathbf{E}\left[\sigma_{S_t,t}^2 S_t^2 \Gamma_{BS}(S_t) | S_0\right]}{\mathbf{E}\left[S_t^2 \Gamma_{BS}(S_t) | S_0\right]}$$
(27)

where we define $\Gamma_{BS}(S_t) := \frac{\partial^2}{\partial S_t^2} C_{BS}(S_t, K, t, T)$.

Now we have a formula for the Black-Scholes implied volatility of a European option in terms of local volatilities. From the definition of $v_{K,T}(t)$, we have that

$$\sigma_{BS}(K,T)^2 = \frac{1}{T} \int_0^T v_{K,T}(t) dt$$

Then, explicitly

$$\sigma_{BS}(K,T)^2 = \frac{1}{T} \int_0^T \frac{\mathbf{E} \left[\sigma_{S_t,t}^2 S_t^2 \Gamma_{BS}(S_t) \left| S_0 \right] \right]}{\mathbf{E} \left[S_t^2 \Gamma_{BS}(S_t) \left| S_0 \right]} dt$$
(28)

Note however that equations (27) and (28) are implicit because the gamma $\Gamma_{BS}(S_t)$ of the option depends on all the forward implied variances $v_{K,T}(t)$.

Special Case (Black-Scholes)

Suppose $\sigma_{S_t,t} = \sigma_t$, a function of t only. Then

$$v_{K,T}(t) = \frac{\mathbf{E}\left[\sigma_t^2 S_t^2 \Gamma_{BS}(S_t) | S_0\right]}{\mathbf{E}\left[S_t^2 \Gamma_{BS}(S_t) | S_0\right]} = \sigma_t^2$$

The forward implied variance $v_{K,T}(t)$ and the local variance σ_t^2 coincide. As expected, $v_{K,T}(t)$ has no dependence on the strike K or the option expiration T.

Interpretation

In order to get better intuition for equation (27), first recall how to compute a risk-neutral expectation:

$$\mathbf{E}\left[f\left(S_{t}\right)\right] = \int dS_{t} p\left(S_{t}, t; S_{0}\right) f\left(S_{t}\right)$$

We get the risk-neutral pdf of the stock price at time t by taking the second derivative of the market price of European options with respect to strike price.

$$p(S_t, t; S_0) = \left. \frac{\partial^2 C(S_0, K, t)}{\partial K^2} \right|_{K=S_t}$$

Then we may rewrite equation (27) as

$$v_{K,T}(t) = \frac{1}{\mathbf{E}[S_t^2 \Gamma_{BS}(S_t) | S_0]} \int dS_t \, p(S_t, t; S_0) \, S_t^2 \Gamma_{BS}(S_t) \, \sigma_{S_t, t}^2$$

= $\int dS_t \, q(S_t, t; S_0, K, T) \, \sigma_{S_t, t}^2$ (29)

where we further define

$$q(S_t, t; S_0, K, T) := \frac{p(S_t, t; S_0) S_t^2 \Gamma_{BS}(S_t)}{\mathbf{E} [S_t^2 \Gamma_{BS}(S_t) | S_0]}$$
(30)

 $q(S_t, t; S_0, K, T)$ is a probability density which looks like a Brownian Bridge density for the stock price given that the initial stock price is S_0 and the time-T stock price is K.

For convenience, we rewrite Equation (29) in terms of $x_t \equiv \log\left(\frac{S_t}{S_0}\right)$. In terms of x_t ,

$$v_{K,T}(t) = \int dx_t q(x_t, t; x_T, T) \sigma_{x_t, t}^2$$
(31)

Figure 1 shows how $q(x_t, t; x_T, T)$ looks in the case of a 1 year European option struck at 1.3 with a flat 20% volatility. We see that $q(x_t, t; x_T, T)$ peaks on a line (which we will denote by \tilde{x}_t) joining the stock price today with the strike price at expiration. Moreover, the density looks roughly symmetric around the peak. This suggests an expansion around the peak \tilde{x}_t (at which the derivative of $q(x_t, t; x_t, T)$ with respect to x_t is zero). Then we write:

$$q(x_t, t; x_T, T) \approx q(\tilde{x}_t, t; x_T, T) + \frac{1}{2} (x_t - \tilde{x}_t)^2 \left. \frac{\partial^2 q}{\partial x_t^2} \right|_{x_t = \tilde{x}_t}$$
(32)

In practice, the local variance $\sigma_{x_t,t}^2$ is typically not so far from linear in x_t in the region where $q(x_t, t; x_T, T)$ is significant so we may further write

$$\sigma_{x_t,t}^2 \approx \sigma_{\tilde{x}_t,t}^2 + (x_t - \tilde{x}_t) \left. \frac{\partial \sigma_{x_t,t}^2}{\partial x_t} \right|_{x_t = \tilde{x}_t}$$
(33)

Substituting (32) and (33) into the integrand in equation (31) gives

$$v_{K,T}(t) \approx \sigma_{\tilde{x}_t,t}^2$$

Figure 1: Graph of the pdf of x_t conditional on $x_T = Log(K)$ for a 1 year European option, strike 1.3 with current stock price = 1 and 20% volatility.



and we may rewrite equation (28) as

$$\sigma_{BS}(K,T)^2 \approx \frac{1}{T} \int_0^T \sigma_{\tilde{x}_{t,t}}^2 dt$$
(34)

In words, equation (39) says that the Black-Scholes implied variance of an option with strike K is given by the integral from valuation date (t = 0) to the expiration date (t = T) of the local variances along the path \tilde{x}_t that maximizes the Brownian Bridge density $q(x_t, t; x_T, T)$.

Of course, in practice, it's not easy to compute the path \tilde{x}_t . However, we now have a very simple picture for the meaning of Black-Scholes implied variance of a European option with a given strike and expiration - it is approximately the integral from today to expiration of local variances along the most probable path for the stock price conditional on the stock price at expiration being the strike price of the option.

6 The Structure of Implied Volatility in the Heston Model

6.1 Local Volatility in the Heston Model

From Section 3.1 with $x_t \equiv Log(S(t)/K)$ and $\mu = 0$, we have

$$dx_t = -\frac{v_t}{2}dt + \sqrt{v_t} \, dZ_t$$

$$dv_t = -\lambda(v_t - \bar{v})dt + \rho\eta\sqrt{v_t} \, dZ_t + \sqrt{1 - \rho^2}\eta\sqrt{v_t} \, dW_t$$
(35)

where dW_t and dZ_t are orthogonal. Eliminating $\sqrt{v_t}dZ_t$, we get

$$dv_t = -\lambda(v_t - \bar{v})dt + \rho\eta \left(dx_t + \frac{1}{2}v_t dt\right) + \sqrt{1 - \rho^2}\eta\sqrt{v_t}dW_t \qquad (36)$$

Our strategy will be to compute local variances in the Heston model and then integrate local variance from valuation date to expiration date to get the BS implied variance following the results of Section 5.

First, consider the unconditional expectation \hat{v}_s of the instantaneous variance at time s. Solving equation (36) gives

$$\hat{v}_s = (v_0 - \bar{v})e^{-\lambda s} + \bar{v}$$

Then define the expected total variance to time t through the relation

$$w_t \equiv \int_0^t \hat{v}_s ds = (v_0 - \bar{v}) \left\{ \frac{1 - e^{-\lambda t}}{\lambda} \right\} + \bar{v} t$$

Finally, let $u_t \equiv \mathbf{E}[v_t | x_T]$ be the expectation of the instantaneous variance at time t conditional on the final value x_T of x.

Ansatz

(By "ansatz", I mean some working assumption which I haven't been able to justify and may not even be true). Without loss of generality, assume $x_0 = 0$. Then,

$$\mathbf{E}[x_s \,|\, x_T] = x_T \frac{w_s}{w_T}$$

where $w_t \equiv \int_0^t ds \, \hat{v}_s$ is the expected total variance to time t. To see that this ansatz is plausible, note that

$$\mathbf{E}(x_s) = \mathbf{E}(x_T)\frac{w_s}{w_T} = -\frac{w_T}{2}\frac{w_s}{w_T} = -\frac{w_s}{2}$$

In fact, if the process for x_t were a conventional Brownian Bridge process, the result would be true but in this case, the result is only approximately true. If you manage to derive the correct result, please let me know.

Assuming the ansatz to be correct, we may take the conditional expectation of (36) to get:

$$du_t = -\lambda(u_t - \bar{v})dt + \frac{\rho\eta}{2}u_t dt + \rho\eta \frac{x_T}{w_T} dw_t + \sqrt{1 - \rho^2} \eta \sqrt{v_t} \mathbf{E}[dW_t | x_T]$$
(37)

If the dependence of dW_t on x_T is weak or if $\sqrt{1-\rho^2}$ is very small, we may drop the last term to get

$$du_t \approx -\lambda'(u_t - \bar{v}')dt + \rho\eta \frac{x_T}{w_T} \hat{v}_t dt$$

with $\lambda' = \lambda - \frac{\rho \eta}{2}$, $\bar{v}' = \bar{v} \frac{\lambda}{\lambda'}$. The solution to this equation is

$$u_T \approx \hat{v}'_T + \rho \eta \, \frac{x_T}{w_T} \, \int_0^T \hat{v}_s \, e^{-\lambda'(T-s)} ds \tag{38}$$

with $\hat{v}'_s \equiv (v - \bar{v}') e^{-\lambda' s} + \bar{v}'$.

From Section 2.5, we know that the local variance $\sigma^2(K, T, S_0) = \mathbf{E} [v_T | S_T = K]$. Then, Equation 38 gives us an approximate but surprisingly accurate formula for local variance within the Heston model (an extremely accurate approximation when $\rho = \pm 1$). We see that in the Heston model, local variance is approximately linear in $x = \log(\frac{F}{K})$.

In summary, we have made two approximations: the Ansatz and dropping the last term in equation (37). For reasonable parameters, equation (38) gives good intuition for the functional form of local variance and when $\rho = \pm 1$, it is almost exact.

6.2 Implied Volatility in the Heston Model

Now, to get implied variance in the Heston model, following the results of Section 5, we need to integrate the Heston local variance along the most probable stock price path joining the initial stock price to the strike price at expiration (the one which maximizes the Brownian Bridge probability density).

In the notation of Section 5, the Black-Scholes implied variance is given by

$$\sigma_{BS}(K,T)^2 \approx \frac{1}{T} \int_0^T \sigma_{\tilde{x}_t,t}^2 dt = \frac{1}{T} \int_0^T u_t(\tilde{x}_t) dt$$
(39)

where $\{\tilde{x}_t\}$ is the most probable path (as defined above).

Recall that the Brownian Bridge density $q(x_t, t; x_T, T)$ is roughly symmetric and peaked around \tilde{x}_t , so $\mathbf{E}[x_t - \tilde{x}_t | x_T] \approx 0$. Applying the Ansatz once again, we obtain

$$\tilde{x}_t = \mathbf{E}\left[\tilde{x}_t | x_T\right] = \mathbf{E}\left[\tilde{x}_t - x_t | x_T\right] + \mathbf{E}\left[x_t | x_T\right] \approx \frac{w_t}{w_T} x_T$$

We substitute this expression back into equations 38 and 39 to get

$$\sigma_{BS}(K,T)^2 \approx \frac{1}{T} \int_0^T u_t(\tilde{x}_t) dt$$
$$\approx \frac{1}{T} \int_0^T \hat{v}_t' dt + \rho \eta \frac{x_T}{w_T} \frac{1}{T} \int_0^T dt \int_0^t \hat{v}_s e^{-\lambda'(t-s)} ds \quad (40)$$

The BS Implied Volatility Term Structure in the Heston Model

The at-the-money term structure of BS implied variance in the Heston model is obtained by setting $x_T = 0$ in equation (40). Performing the integration explicitly gives

$$\sigma_{BS}(K,T)^{2}\Big|_{K=F_{T}} \approx \frac{1}{T} \int_{0}^{T} \hat{v}_{t}' dt = \frac{1}{T} \int_{0}^{T} \left[(v - \bar{v}') e^{-\lambda' t} + \bar{v}' \right] dt$$
$$= (v - \bar{v}') \frac{1 - e^{-\lambda' T}}{\lambda' T} + \bar{v}'$$

We see that in the Heston model, the at-the-money Black-Scholes implied variance $\sigma_{BS}(K,T)^2|_{K=F_T} \to v$ (the instantaneous variance) as the time to expiration $T \to 0$ and as $T \to \infty$, the at-the-money Black-Scholes implied variance reverts to \bar{v}' .

The BS Implied Volatility Skew in the Heston Model

It is possible (but not very illuminating) to integrate the second term of equation (40) explicitly. Even without doing that, we can see that the implied variance skew in the Heston model is approximately linear in the correlation ρ and the volatility of volatility η .

In the special case where $v_0 = \bar{v}$, the implied variance skew has a particularly simple form. Then $\hat{v}_s = \bar{v}$ and $w_t = \bar{v}t$. The most probable path $\tilde{x}_t \approx \frac{t}{T}x_T$ is exactly a straight line in log-space between the initial stock price on valuation date and the strike price at expiration. Performing the integrations in Equation 40 explicitly, we get

$$\sigma_{BS}(K,T)^2 \approx \frac{w_T'}{T} + \rho \eta \frac{x_T}{T^2} \int_0^T dt \frac{1}{T} \int_0^t e^{-\lambda'(t-s)} ds$$
$$= \frac{w_T'}{T} + \rho \eta \frac{x_T}{\lambda'T} \left\{ 1 - \frac{\left(1 - e^{-\lambda'T}\right)}{\lambda'T} \right\}$$
(41)

From equation (41), we see that the implied variance skew $\frac{\partial}{\partial x_t} \sigma_{BS}(K,T)^2$ is *independent* of the level of instantaneous variance v or long-term mean variance \bar{v} . In fact, this remains approximately true even when $v \neq \bar{v}$. It follows that we now have a fast way of calibrating the Heston model to observed implied volatility skews. Just two expirations would in principle allow us to determine λ' and the product $\rho\eta$. We can then fit the term structure of volatility to determine the long term mean variance \bar{v} and the instantaneous variance v_0 . The curvature of the skew (not discussed here) would allow us to determine ρ and η separately.

We note that as we increase either the correlation ρ or the volatility of volatility η , the skew increases.

Also, the very short-dated skew is independent of λ and T:

$$\frac{\partial}{\partial x_t} \sigma_{BS}(K,T)^2 = \rho \eta \frac{1}{\lambda' T} \left\{ 1 - \frac{\left(1 - e^{-\lambda' T}\right)}{\lambda' T} \right\} \to \frac{\rho \eta}{2} \quad \text{as} \quad T \to 0$$

and the long-dated skew is inversely proportional to T:

$$\frac{\partial}{\partial x_t} \sigma_{BS}(K,T)^2 = \rho \eta \frac{1}{\lambda' T} \left\{ 1 - \frac{\left(1 - e^{-\lambda' T}\right)}{\lambda' T} \right\} \sim \frac{\rho \eta}{\lambda' T} \quad \text{as} \quad T \to \infty$$

Finally, increasing η causes the curvature of the implied volatility skew (related to the kurtosis of the risk-neutral density) to increase but we haven't shown that here.

7 The SPX Implied Volatility Surface

Up to this point, we have concentrated on understanding the shape of the implied volatility surface implied by a stochastic volatility model – in particular the Heston model. However, we still have no idea whether implied volatilities produced by the Heston model look like implied volatilities in the market. In the accompanying file (SPXvols000128.pdf), you can find graphs of the SPX implied volatility surface as of January 28, 2000.

After fitting a quadratic to observed variance as a function of x_t for each expiration t, we get the at-the-money forward variance levels and skews listed in Table 1. (Recall that by at-the-money skew, I mean $\frac{\partial}{\partial x_t}\sigma_{BS}(K,T)^2$).

Expiration	Time (years)	ATM Variance	ATM Skew
17-Feb-2000	0.0548	0.0619	-0.2586
16-Mar-2000	0.1315	0.0550	-0.2360
20-Apr-2000	0.2274	0.0528	-0.2013
15-Jun-2000	0.3808	0.0532	-0.1722
14-Sep-2000	0.6301	0.0539	-0.1453
14-Dec-2000	0.8795	0.0560	-0.1292
14-Jun-2001	1.3781	0.0574	-0.1108
20-Dec-2001	1.8959	0.0554	-0.1018

Table 1: At-the-money SPX variance levels and skews on January 28, 2000.

Skew is plotted as a function of time in Figure 2. Just looking at the pattern of the points, we would suspect that a simple functional form should be able to fit. However, the solid and dashed lines show the results of fitting the approximate formula

$$\rho\eta\frac{1}{\lambda'T}\left\{1-\frac{\left(1-e^{-\lambda'T}\right)}{\lambda'T}\right\}$$

to the observed skews. The solid line takes all points into account; the dashed line drops the first three expirations from the fit. We can see that

the fitting function is too stiff to fit the observed pattern of variance skews; there is no choice of λ' that will allow us to fit the skew observations. The fact that the observed variance skew increases significantly faster as $T \rightarrow 0$ than the skew implied by a stochastic volatility model may indicate that jumps need to be included in a complete model as in Matytsin (1999) for example.

Figure 2: Graph of SPX ATM skew vs. time. The dashed fit excludes the first 3 data points.

In Figure 3, we see that trying to fit at-the-money variance using a simple model is a pretty hopeless endeavor. Once again, for reference, we show the solid and dashed line best fits of the function

$$\sigma_{BS}(K,T)^2\Big|_{K=F_T} \approx (v-\bar{v}') \frac{1}{\lambda'T} \left\{ 1 - \frac{\left(1-e^{-\lambda'T}\right)}{\lambda'T} \right\} + \bar{v}'$$

Now we understand one reason why practitioners prefer local volatility models – a stochastic volatility model with time-homogeneous parameters cannot fit market prices! Perhaps an extended stochastic volatility model with correlated jumps in stock price and volatility (Matytsin (1999)) might fit better? But how would traders choose their input parameters? How would the SPX index book trader choose his volatility of volatility parameter – or worse, the correlation between jumps in stock price and jumps in volatility ?

Figure 3: Graph of SPX ATM variance vs. time. The dashed fit excludes the first 3 data points.

References

- Avellaneda, Marco, C. Friedman, R. Holmes, and D. Samperi, 1997, Calibrating volatility surfaces via relative-entropy minimization, *Applied Mathematical Finance* 4, 37–64.
- Blacher, Guillaume, 1998, Local volatility, RISK Conference Presentation.
- Derman, Emanuel, Iraj Kani, and Neil Chriss, 1996, Implied trinomial trees of the volatility smile, *Journal of Derivatives* 3, 7–22.
- Jackwerth, Jens C., 2000, Option-implied risk-neutral distributions and implied binomial trees: A literature review, *Journal of Derivatives* 7, 66–82.
- Matytsin, Andrew, 1999, Modelling volatility and volatility derivatives, Columbia Practitioners Conference on the Mathematics of Finance.
- Rubinstein, Mark, 1998, Edgeworth binomial trees, *Journal of Derivatives* 5, 20–27.
- Shimko, D., 1993, Bounds on probability, Risk 6, 33–37.